

THE POLAR FORM OF COMPLEX NUMBERS

In this section, we return to our study of complex numbers which were first introduced in Section ?? . Recall that a **complex number** is a number of the form $z = a + bi$ where a and b are real numbers and i is the imaginary unit defined by $i = \sqrt{-1}$.

The number a is called the **real part** of z , denoted $\operatorname{Re}(z)$, while the real number b is called the **imaginary part** of z , denoted $\operatorname{Im}(z)$. From Intermediate Algebra, we know that if $z = a + bi = c + di$ where a, b, c and d are real numbers, then $a = c$ and $b = d$, which means $\operatorname{Re}(z)$ and $\operatorname{Im}(z)$ are well-defined.¹

To start off this section, we associate each complex number $z = a + bi$ with the point (a, b) on the Cartesian (rectangular) coordinate plane. In this case, the x -axis is relabeled as the **real axis**, which corresponds to the real number line as usual, and the y -axis is relabeled as the **imaginary axis**, which is demarcated in increments of the imaginary unit i . The plane determined by these two axes is called the **complex plane**.



The Complex Plane

Since the ordered pair (a, b) gives the *rectangular* coordinates associated with $z = a + bi$, the expression $z = a + bi$ is called the **rectangular form** of the complex number z .

We could just as easily associate z with a pair of *polar* coordinates (r, θ) . Although it is not as straightforward as the definitions of $\operatorname{Re}(z)$ and $\operatorname{Im}(z)$, we give r and θ special names in relation to z below.

Definition 1. The Modulus and Argument of Complex Numbers:

Let $z = a + bi$ be a complex number with $a = \operatorname{Re}(z)$ and $b = \operatorname{Im}(z)$. Let (r, θ) be a polar representation of the point with rectangular coordinates (a, b) where $r \geq 0$.

- The **modulus** of z , denoted $|z|$, is defined by $|z| = r$.
- The angle θ is an **argument** of z . The set of all arguments of z is denoted $\arg(z)$.
- If $z \neq 0$ and $-\pi < \theta \leq \pi$, then θ is the **principal argument** of z , written $\theta = \operatorname{Arg}(z)$.

Some remarks about Definition 1 are in order. We know from Section ?? that every point in the plane has infinitely many polar coordinate representations (r, θ) which means it's worth our time to make sure the quantities 'modulus', 'argument' and 'principal argument' are well-defined.

Concerning the modulus, if $z = 0$ then the point associated with z is the origin. In this case, the *only* r -value which can be used here is $r = 0$. Hence for $z = 0$, $|z| = 0$ is well-defined.

If $z \neq 0$, then the point associated with z is not the origin, and there are *two* possibilities for r : one positive and one negative. However, we stipulated $r \geq 0$ in our definition so this pins down the value of $|z|$ to one and only one number. Thus the modulus is well-defined in this case, too.²

¹'Well-defined' means that no matter how we express z , the number $\operatorname{Re}(z)$ is always the same, and the number $\operatorname{Im}(z)$ is always the same. In other words, Re and Im are *functions* of complex numbers.

²In case you're wondering, the use of the absolute value notation $|z|$ for modulus will be explained shortly.

THE POLAR FORM OF COMPLEX NUMBERS

Even with the requirement $r \geq 0$, there are infinitely many angles θ which can be used in a polar representation of a point (r, θ) . If $z \neq 0$ then the point in question is not the origin, so all of these angles θ are coterminal. Since coterminal angles are exactly 2π radians apart, we are guaranteed that only one of them lies in the interval $(-\pi, \pi]$, and this angle is what we call the principal argument of z , $\text{Arg}(z)$.

The set $\arg(z)$ of all arguments of z can be described as $\arg(z) = \{\text{Arg}(z) + 2\pi k \mid k \text{ is an integer}\}$. Note that since $\arg(z)$ is a set, we will write ' $\theta \in \arg(z)$ ' to mean ' θ is in³ the set of arguments of z '.

If $z = 0$ then the point in question is the origin, which we know can be represented in polar coordinates as $(0, \theta)$ for any angle θ . In this case, we have $\arg(0) = (-\infty, \infty)$ and since there is no one value of θ which lies $(-\pi, \pi]$, we leave $\text{Arg}(0)$ undefined. It is time for an example.

Example 1. For each of the following complex numbers find $\text{Re}(z)$, $\text{Im}(z)$, $|z|$, $\arg(z)$ and $\text{Arg}(z)$. Plot each complex number z in the complex plane.

1. $z = \sqrt{3} - i$

2. $z = -2 + 4i$

3. $z = 3i$

4. $z = -117$

Solution.

1. For $z = \sqrt{3} - i = \sqrt{3} + (-1)i$, we have $\text{Re}(z) = \sqrt{3}$ and $\text{Im}(z) = -1$. To find $|z|$, $\arg(z)$ and $\text{Arg}(z)$, we need to find a polar representation (r, θ) with $r \geq 0$ for the point $P(\sqrt{3}, -1)$ associated with z .

We know $r^2 = (\sqrt{3})^2 + (-1)^2 = 4$, so $r = \pm 2$. Since we require $r \geq 0$, we choose $r = 2$, so $|z| = 2$.

To find a corresponding angle θ , we note that since $r > 0$ and P lies in Quadrant IV, θ must be a Quadrant IV angle. We know $\tan(\theta) = \frac{-1}{\sqrt{3}} = -\frac{\sqrt{3}}{3}$, so $\theta = -\frac{\pi}{6} + 2\pi k$ for integers k . Hence, $\arg(z) = \left\{-\frac{\pi}{6} + 2\pi k \mid k \text{ is an integer}\right\}$. Of these values, only $\theta = -\frac{\pi}{6}$ satisfies $-\pi < \theta \leq \pi$, hence we get $\text{Arg}(z) = -\frac{\pi}{6}$.

2. The complex number $z = -2 + 4i$ has $\text{Re}(z) = -2$, $\text{Im}(z) = 4$, and is associated with the point $P(-2, 4)$. Our next task is to find a polar representation (r, θ) for P where $r \geq 0$.

Running through the usual calculations gives $r = 2\sqrt{5}$, so $|z| = 2\sqrt{5}$. To find θ , we get $\tan(\theta) = -2$, and since $r > 0$ and P lies in Quadrant II, we know θ is a Quadrant II angle.

We find $\theta = \pi + \arctan(-2) + 2\pi k$, or, more succinctly $\theta = \pi - \arctan(2) + 2\pi k$ for integers k . Hence $\arg(z) = \{\pi - \arctan(2) + 2\pi k \mid k \text{ is an integer}\}$. Only $\theta = \pi - \arctan(2)$ satisfies $-\pi < \theta \leq \pi$, so we get $\text{Arg}(z) = \pi - \arctan(2)$.

3. We rewrite $z = 3i$ as $z = 0 + 3i$ to find $\text{Re}(z) = 0$ and $\text{Im}(z) = 3$. The point in the plane which corresponds to z is $(0, 3)$ and while we could go through the usual calculations to find the required polar form of this point, we can obtain the answer 'by inspection.'

The point $(0, 3)$ lies 3 units away from the origin on the positive y -axis. Hence, $r = |z| = 3$ and $\theta = \frac{\pi}{2} + 2\pi k$ for integers k . We get $\arg(z) = \left\{\frac{\pi}{2} + 2\pi k \mid k \text{ is an integer}\right\}$ and $\text{Arg}(z) = \frac{\pi}{2}$.

4. As in the previous problem, we write $z = -117 = -117 + 0i$ so $\text{Re}(z) = -117$ and $\text{Im}(z) = 0$. The number $z = -117$ corresponds to the point $(-117, 0)$, and this is another instance where we can determine the polar form 'by eye'.

The point $(-117, 0)$ is 117 units away from the origin along the negative x -axis. Hence, $r = |z| = 117$ and $\theta = \pi + 2\pi k = (2k + 1)\pi$ for integers k . We have $\arg(z) = \{(2k + 1)\pi \mid k \text{ is an integer}\}$.

Only one of these values, $\theta = \pi$, (just barely!) lies in the interval $(-\pi, \pi]$ which means $\text{Arg}(z) = \pi$. We plot z along with the other numbers in this example below.

³Recall the symbol being used here, ' $\in$,' is the mathematical symbol which denotes membership in a set. See Section ??.

THE POLAR FORM OF COMPLEX NUMBERS



Now that we've had practice computing the modulus of a complex number, we state some properties below.

Theorem 1. Properties of the Modulus: Let z and w be complex numbers.

- $|z|$ is the distance from z to 0 in the complex plane
- $|z| \geq 0$ and $|z| = 0$ if and only if $z = 0$
- $|z| = \sqrt{\operatorname{Re}(z)^2 + \operatorname{Im}(z)^2}$
- **Product Rule:** $|zw| = |z||w|$
- **Power Rule:** $|z^n| = |z|^n$ for all natural numbers, n
- **Quotient Rule:** $\left| \frac{z}{w} \right| = \frac{|z|}{|w|}$, provided $w \neq 0$

To prove the first three properties in Theorem 1, suppose $z = a + bi$ where a and b are real numbers. To determine $|z|$, we find a polar representation (r, θ) with $r \geq 0$ for the point (a, b) .

From Section ??, we know $r^2 = a^2 + b^2$ so that $r = \pm\sqrt{a^2 + b^2}$. Since we require $r \geq 0$, then it must be that $r = \sqrt{a^2 + b^2}$, which means $|z| = \sqrt{a^2 + b^2}$. Using the distance formula, we find the distance from $(0, 0)$ to (a, b) is also $\sqrt{a^2 + b^2}$, establishing the first property.⁴

For the second property, note that since $|z|$ is a distance, $|z| \geq 0$. Furthermore, $|z| = 0$ if and only if the distance from z to 0 is 0, and the latter happens if and only if $z = 0$, which is what we were asked to show.⁵

For the third property, we note that since $a = \operatorname{Re}(z)$ and $b = \operatorname{Im}(z)$, $z = \sqrt{a^2 + b^2} = \sqrt{\operatorname{Re}(z)^2 + \operatorname{Im}(z)^2}$.

To prove the product rule, suppose $z = a + bi$ and $w = c + di$ for real numbers a, b, c and d . Then $zw = (a + bi)(c + di)$. After the usual arithmetic⁶ we get $zw = (ac - bd) + (ad + bc)i$. Therefore,

$$\begin{aligned}
 |zw| &= \sqrt{(ac - bd)^2 + (ad + bc)^2} \\
 &= \sqrt{a^2c^2 - 2abcd + b^2d^2 + a^2d^2 + 2abcd + b^2c^2} && \text{Expand} \\
 &= \sqrt{a^2c^2 + a^2d^2 + b^2c^2 + b^2d^2} && \text{Rearrange terms} \\
 &= \sqrt{a^2(c^2 + d^2) + b^2(c^2 + d^2)} && \text{Factor} \\
 &= \sqrt{(a^2 + b^2)(c^2 + d^2)} && \text{Factor} \\
 &= \sqrt{a^2 + b^2} \sqrt{c^2 + d^2} && \text{Product Rule for Radicals} \\
 &= |z||w| && \text{Definition of } |z| \text{ and } |w|
 \end{aligned}$$

⁴Since the absolute value $|x|$ of a real number x can be viewed as the distance from x to 0 on the number line, this first property justifies the notation $|z|$ for modulus. We leave it to the reader to show that if z is real, then the definition of modulus coincides with absolute value so the notation $|z|$ is unambiguous.

⁵This may be considered by some to be a bit of a cheat, so we work through the underlying Algebra to see this is true. We know $|z| = 0$ if and only if $\sqrt{a^2 + b^2} = 0$ if and only if $a^2 + b^2 = 0$, which is true if and only if $a = b = 0$. The latter happens if and only if $z = a + bi = 0$. There.

⁶See Example ?? in Section ?? for a review of complex number arithmetic.

THE POLAR FORM OF COMPLEX NUMBERS

Hence $|zw| = |z||w|$ as required.

Now that the Product Rule has been established, we use it and the Principle of Mathematical Induction⁷ to prove the power rule. Let $P(n)$ be the statement $|z^n| = |z|^n$. Then $P(1)$ is true since $|z^1| = |z| = |z|^1$.

Next, assume $P(k)$ is true. That is, assume $|z^k| = |z|^k$ for some $k \geq 1$. Our job is to show that $P(k+1)$ is true, namely $|z^{k+1}| = |z|^{k+1}$. As is customary with induction proofs, we first try to reduce the problem in such a way as to use the Induction Hypothesis.

$$\begin{aligned} |z^{k+1}| &= |z^k z| && \text{Properties of Exponents} \\ &= |z^k| |z| && \text{Product Rule} \\ &= |z|^k |z| && \text{Induction Hypothesis} \\ &= |z|^{k+1} && \text{Properties of Exponents} \end{aligned}$$

Hence, $P(k+1)$ is true, which means $|z^n| = |z|^n$ is true for all natural numbers n .

Like the Power Rule, the Quotient Rule can also be established with the help of the Product Rule. We assume $w \neq 0$ (so $|w| \neq 0$) and we get

$$\begin{aligned} \left| \frac{z}{w} \right| &= \left| \left(z \right) \left(\frac{1}{w} \right) \right| \\ &= |z| \left| \frac{1}{w} \right| && \text{Product Rule.} \end{aligned}$$

Hence, the proof really boils down to showing $\left| \frac{1}{w} \right| = \frac{1}{|w|}$. This is left as an exercise.

Next, we characterize the argument of a complex number in terms of its real and imaginary parts.

Theorem 2. Properties of the Argument: Let z be a complex number.

- If $\operatorname{Re}(z) \neq 0$ and $\theta \in \arg(z)$, then $\tan(\theta) = \frac{\operatorname{Im}(z)}{\operatorname{Re}(z)}$.
- If $\operatorname{Re}(z) = 0$ and $\operatorname{Im}(z) > 0$, then $\arg(z) = \left\{ \frac{\pi}{2} + 2\pi k \mid k \text{ is an integer} \right\}$.
- If $\operatorname{Re}(z) = 0$ and $\operatorname{Im}(z) < 0$, then $\arg(z) = \left\{ -\frac{\pi}{2} + 2\pi k \mid k \text{ is an integer} \right\}$.
- If $\operatorname{Re}(z) = \operatorname{Im}(z) = 0$, then $z = 0$ and $\arg(z) = (-\infty, \infty)$.

To prove Theorem 2, suppose $z = a + bi$ for real numbers a and b . By definition, $a = \operatorname{Re}(z)$ and $b = \operatorname{Im}(z)$, so the point associated with z is $(a, b) = (\operatorname{Re}(z), \operatorname{Im}(z))$. From Section ??, we know that if (r, θ) is a polar representation for $(\operatorname{Re}(z), \operatorname{Im}(z))$, then $\tan(\theta) = \frac{\operatorname{Im}(z)}{\operatorname{Re}(z)}$, provided $\operatorname{Re}(z) \neq 0$.

If $\operatorname{Re}(z) = 0$ and $\operatorname{Im}(z) > 0$, then z lies on the positive imaginary axis. Since we take $r > 0$, we have that θ is coterminal with $\frac{\pi}{2}$, and the result follows. If $\operatorname{Re}(z) = 0$ and $\operatorname{Im}(z) < 0$, then z lies on the negative imaginary axis, and a similar argument shows θ is coterminal with $-\frac{\pi}{2}$.

The last property in the theorem was already discussed in the remarks following Definition 1.

Our next goal is to completely marry the Geometry and the Algebra of the complex numbers. To that end, consider the figure below.

⁷See Section ?? for a review of this technique.

THE POLAR FORM OF COMPLEX NUMBERS



Polar coordinates, (r, θ) associated with $z = a + bi$ with $r \geq 0$.

We know from Theorem ?? that $a = r \cos(\theta)$ and $b = r \sin(\theta)$. Making these substitutions for a and b gives $z = a + bi = r \cos(\theta) + r \sin(\theta)i = r[\cos(\theta) + i \sin(\theta)]$.

The expression ' $\cos(\theta) + i \sin(\theta)$ ' is abbreviated $\text{cis}(\theta)$ so we can write $z = r \text{cis}(\theta) = |z| \text{cis}(\theta)$.

Definition 2. A Polar Form of a Complex Number:

Suppose z is a complex number and $\theta \in \arg(z)$. The expression:

$$|z| \text{cis}(\theta) = |z| [\cos(\theta) + i \sin(\theta)]$$

is called a polar form for z .

Since there are infinitely many choices for $\theta \in \arg(z)$, there are infinitely many polar forms for z , so we used the indefinite article 'a' in Definition 2. It is time for an example.

Example 2.

- Find the rectangular form of the following complex numbers. Find $\text{Re}(z)$ and $\text{Im}(z)$.

$$(i) \ z = 4 \text{cis} \left(\frac{2\pi}{3} \right) \quad (ii) \ z = 2 \text{cis} \left(-\frac{3\pi}{4} \right) \quad (iii) \ z = 3 \text{cis}(0) \quad (iv) \ z = \text{cis} \left(\frac{\pi}{2} \right)$$

- Use the results from Example 1 to find a polar form of the following complex numbers.

$$(i) \ z = \sqrt{3} - i \quad (ii) \ z = -2 + 4i \quad (iii) \ z = 3i \quad (iv) \ z = -117$$

Solution.

- The key to this problem is to write out $\text{cis}(\theta)$ as $\cos(\theta) + i \sin(\theta)$.

- By definition, $z = 4 \text{cis} \left(\frac{2\pi}{3} \right) = 4 \left[\cos \left(\frac{2\pi}{3} \right) + i \sin \left(\frac{2\pi}{3} \right) \right]$. Simplifying, we get $z = -2 + 2i\sqrt{3}$, so that $\text{Re}(z) = -2$ and $\text{Im}(z) = 2\sqrt{3}$.
- Expanding, we get $z = 2 \text{cis} \left(-\frac{3\pi}{4} \right) = 2 \left[\cos \left(-\frac{3\pi}{4} \right) + i \sin \left(-\frac{3\pi}{4} \right) \right]$. Hence, $z = -\sqrt{2} - i\sqrt{2}$, so $\text{Re}(z) = -\sqrt{2} = \text{Im}(z)$.
- We get $z = 3 \text{cis}(0) = 3[\cos(0) + i \sin(0)] = 3$. Writing $3 = 3 + 0i$, we get $\text{Re}(z) = 3$ and $\text{Im}(z) = 0$, which makes sense seeing as 3 is a real number.
- Lastly, we have $z = \text{cis} \left(\frac{\pi}{2} \right) = \cos \left(\frac{\pi}{2} \right) + i \sin \left(\frac{\pi}{2} \right) = i$. Since $i = 0 + 1i$, we get $\text{Re}(z) = 0$ and $\text{Im}(z) = 1$. Since i is called the 'imaginary unit,' these answers make perfect sense.

THE POLAR FORM OF COMPLEX NUMBERS

2. To write a polar form of a complex number z , we need two pieces of information: the modulus $|z|$ and an argument (not necessarily the principal argument) of z .

We shamelessly mine our solution to Example 1 to find what we need.

- (i) For $z = \sqrt{3} - i$, $|z| = 2$ and $\theta = -\frac{\pi}{6}$, so $z = 2\text{cis}\left(-\frac{\pi}{6}\right)$. We can check our answer by converting it back to rectangular form to see that it simplifies to $z = \sqrt{3} - i$.
- (ii) For $z = -2 + 4i$, $|z| = 2\sqrt{5}$ and $\theta = \pi - \arctan(2)$. Hence, $z = 2\sqrt{5}\text{cis}(\pi - \arctan(2))$. It is a good exercise to actually show that this polar form reduces to $z = -2 + 4i$.
- (iii) For $z = 3i$, $|z| = 3$ and $\theta = \frac{\pi}{2}$. In this case, $z = 3\text{cis}\left(\frac{\pi}{2}\right)$. This can be checked geometrically.
Head out 3 units from 0 along the positive real axis. Rotating $\frac{\pi}{2}$ radians counter-clockwise lands you exactly 3 units above 0 on the imaginary axis at $z = 3i$.
- (iv) Last but not least, for $z = -117$, $|z| = 117$ and $\theta = \pi$. We get $z = 117\text{cis}(\pi)$. As with the previous problem, our answer is easily checked geometrically. ■

The following theorem summarizes the advantages of working with complex numbers in polar form.

Theorem 3. Products, Powers and Quotients Complex Numbers in Polar Form:

Suppose z and w are complex numbers with polar forms $z = |z|\text{cis}(\alpha)$ and $w = |w|\text{cis}(\beta)$. Then

- **Product Rule:** $zw = |z||w|\text{cis}(\alpha + \beta)$
- **Power Rule (DeMoivre's Theorem) :** $z^n = |z|^n\text{cis}(n\theta)$ for every natural number n
- **Quotient Rule:** $\frac{z}{w} = \frac{|z|}{|w|}\text{cis}(\alpha - \beta)$, provided $|w| \neq 0$

The proof of Theorem 3 requires a healthy mix of definition, arithmetic and identities. We first start with the product rule.

$$\begin{aligned} zw &= [|z|\text{cis}(\alpha)] [|w|\text{cis}(\beta)] \\ &= |z||w| [\cos(\alpha) + i\sin(\alpha)] [\cos(\beta) + i\sin(\beta)] \end{aligned}$$

We now focus on the quantity in brackets on the right hand side of the equation.

$$\begin{aligned} [\cos(\alpha) + i\sin(\alpha)] [\cos(\beta) + i\sin(\beta)] &= \cos(\alpha)\cos(\beta) + i\cos(\alpha)\sin(\beta) \\ &\quad + i\sin(\alpha)\cos(\beta) + i^2\sin(\alpha)\sin(\beta) \\ &= \cos(\alpha)\cos(\beta) + i^2\sin(\alpha)\sin(\beta) && \text{Rearranging terms} \\ &\quad + i\sin(\alpha)\cos(\beta) + i\cos(\alpha)\sin(\beta) \\ &= (\cos(\alpha)\cos(\beta) - \sin(\alpha)\sin(\beta)) && \text{Since } i^2 = -1 \\ &\quad + i(\sin(\alpha)\cos(\beta) + \cos(\alpha)\sin(\beta)) && \text{Factor out } i \\ &= \cos(\alpha + \beta) + i\sin(\alpha + \beta) && \text{Sum identities} \\ &= \text{cis}(\alpha + \beta) && \text{Definition of 'cis'} \end{aligned}$$

Putting this together with our earlier work, we get $zw = |z||w|\text{cis}(\alpha + \beta)$, as required.

Next take aim at the Power Rule, better known as DeMoivre's Theorem.⁸ We proceed by induction on n . Let $P(n)$ be the sentence $z^n = |z|^n\text{cis}(n\theta)$. Then $P(1)$ is true, since $z^1 = z = |z|\text{cis}(\theta) = |z|^1\text{cis}(1 \cdot \theta)$.

⁸Compare this proof with the proof of the Power Rule in Theorem 1.

THE POLAR FORM OF COMPLEX NUMBERS

We now assume $P(k)$ is true, that is, we assume $z^k = |z|^k \text{cis}(k\theta)$ for some $k \geq 1$. Our goal is to show that $P(k+1)$ is true, or that $z^{k+1} = |z|^{k+1} \text{cis}((k+1)\theta)$. We have

$$\begin{aligned} z^{k+1} &= z^k z && \text{Properties of Exponents} \\ &= (|z|^k \text{cis}(k\theta)) (|z| \text{cis}(\theta)) && \text{Induction Hypothesis} \\ &= (|z|^k |z|) \text{cis}(k\theta + \theta) && \text{Product Rule} \\ &= |z|^{k+1} \text{cis}((k+1)\theta) \end{aligned}$$

Hence, assuming $P(k)$ is true, we have that $P(k+1)$ is true, so by the Principle of Mathematical Induction, $z^n = |z|^n \text{cis}(n\theta)$ for all natural numbers n .

The last property in Theorem 3 to prove is the quotient rule. Assuming $|w| \neq 0$ we have

$$\begin{aligned} \frac{z}{w} &= \frac{|z| \text{cis}(\alpha)}{|w| \text{cis}(\beta)} \\ &= \left(\frac{|z|}{|w|} \right) \frac{\cos(\alpha) + i \sin(\alpha)}{\cos(\beta) + i \sin(\beta)} \end{aligned}$$

Next, we multiply both the numerator and denominator of the right hand side by $(\cos(\beta) - i \sin(\beta))$ which is the complex conjugate of $(\cos(\beta) + i \sin(\beta))$ to get

$$\frac{z}{w} = \left(\frac{|z|}{|w|} \right) \frac{\cos(\alpha) + i \sin(\alpha)}{\cos(\beta) + i \sin(\beta)} \cdot \frac{\cos(\beta) - i \sin(\beta)}{\cos(\beta) - i \sin(\beta)}$$

If we let the numerator be $N = [\cos(\alpha) + i \sin(\alpha)][\cos(\beta) - i \sin(\beta)]$ and simplify we get

$$\begin{aligned} N &= [\cos(\alpha) + i \sin(\alpha)][\cos(\beta) - i \sin(\beta)] \\ &= \cos(\alpha) \cos(\beta) - i \cos(\alpha) \sin(\beta) + i \sin(\alpha) \cos(\beta) - i^2 \sin(\alpha) \sin(\beta) && \text{Expand} \\ &= [\cos(\alpha) \cos(\beta) + \sin(\alpha) \sin(\beta)] + i [\sin(\alpha) \cos(\beta) - \cos(\alpha) \sin(\beta)] && \text{Rearrange and Factor} \\ &= \cos(\alpha - \beta) + i \sin(\alpha - \beta) && \text{Difference Identities} \\ &= \text{cis}(\alpha - \beta) && \text{Definition of 'cis'} \end{aligned}$$

If we call the denominator D then we get

$$\begin{aligned} D &= [\cos(\beta) + i \sin(\beta)][\cos(\beta) - i \sin(\beta)] \\ &= \cos^2(\beta) - i \cos(\beta) \sin(\beta) + i \cos(\beta) \sin(\beta) - i^2 \sin^2(\beta) && \text{Expand} \\ &= \cos^2(\beta) - i^2 \sin^2(\beta) && \text{Simplify} \\ &= \cos^2(\beta) + \sin^2(\beta) && \text{Again, } i^2 = -1 \\ &= 1 && \text{Pythagorean Identity} \end{aligned}$$

Putting it all together, we get

$$\begin{aligned} \frac{z}{w} &= \left(\frac{|z|}{|w|} \right) \frac{\cos(\alpha) + i \sin(\alpha)}{\cos(\beta) + i \sin(\beta)} \cdot \frac{\cos(\beta) - i \sin(\beta)}{\cos(\beta) - i \sin(\beta)} \\ &= \left(\frac{|z|}{|w|} \right) \frac{\text{cis}(\alpha - \beta)}{1} \\ &= \frac{|z|}{|w|} \text{cis}(\alpha - \beta) \end{aligned}$$

and we are done. The next example makes good use of Theorem 3.

Example 3. Let $z = 2\sqrt{3} + 2i$ and $w = -1 + i\sqrt{3}$. Use Theorem 3 to find the following.

THE POLAR FORM OF COMPLEX NUMBERS

1. zw

2. w^5

3. $\frac{z}{w}$

Write your final answers in rectangular form.

Solution. In order to use Theorem 3, we need to write z and w in polar form.

For $z = 2\sqrt{3} + 2i$, we find $|z| = \sqrt{(2\sqrt{3})^2 + (2)^2} = \sqrt{16} = 4$. If $\theta \in \arg(z)$, then $\tan(\theta) = \frac{\operatorname{Im}(z)}{\operatorname{Re}(z)} = \frac{2}{2\sqrt{3}} = \frac{\sqrt{3}}{3}$.

Since z lies in Quadrant I, we have $\theta = \frac{\pi}{6} + 2\pi k$ for integers k . Hence, $z = 4\operatorname{cis}\left(\frac{\pi}{6}\right)$.

For $w = -1 + i\sqrt{3}$, we have $|w| = \sqrt{(-1)^2 + (\sqrt{3})^2} = 2$. For an argument θ of w , $\tan(\theta) = \frac{\sqrt{3}}{-1} = -\sqrt{3}$. Since

w lies in Quadrant II, $\theta = \frac{2\pi}{3} + 2\pi k$ for integers k and $w = 2\operatorname{cis}\left(\frac{2\pi}{3}\right)$.

Since we now have polar forms of z and w , we can now proceed using Theorem 3.

$$1. \text{ We get } zw = \left(4\operatorname{cis}\left(\frac{\pi}{6}\right)\right) \left(2\operatorname{cis}\left(\frac{2\pi}{3}\right)\right) = 8\operatorname{cis}\left(\frac{\pi}{6} + \frac{2\pi}{3}\right) = 8\operatorname{cis}\left(\frac{5\pi}{6}\right) = 8 \left[\cos\left(\frac{5\pi}{6}\right) + i \sin\left(\frac{5\pi}{6}\right) \right].$$

After simplifying, we get $zw = -4\sqrt{3} + 4i$.

$$2. \text{ We use DeMoivre's Theorem which yields } w^5 = \left[2\operatorname{cis}\left(\frac{2\pi}{3}\right)\right]^5 = 2^5\operatorname{cis}\left(5 \cdot \frac{2\pi}{3}\right) = 32\operatorname{cis}\left(\frac{10\pi}{3}\right).$$

Since $\frac{10\pi}{3}$ is coterminal with $\frac{4\pi}{3}$, we get $w^5 = 32 \left[\cos\left(\frac{4\pi}{3}\right) + i \sin\left(\frac{4\pi}{3}\right) \right] = -16 - 16i\sqrt{3}$.

$$3. \text{ Last, but not least, we have } \frac{z}{w} = \frac{4\operatorname{cis}\left(\frac{\pi}{6}\right)}{2\operatorname{cis}\left(\frac{2\pi}{3}\right)} = \frac{4}{2}\operatorname{cis}\left(\frac{\pi}{6} - \frac{2\pi}{3}\right) = 2\operatorname{cis}\left(-\frac{\pi}{2}\right).$$

Since $-\frac{\pi}{2}$ is a quadrantal angle, we can 'see' the rectangular form by moving out 2 units along the positive real axis, then rotating $\frac{\pi}{2}$ radians *clockwise* to arrive at the point 2 units below 0 on the imaginary axis. The long and short of it is that $\frac{z}{w} = -2i$. ■

Some remarks are in order. First, the reader may not be sold on using the polar form of complex numbers to multiply complex numbers – especially if they aren't given in polar form to begin with.

Indeed, a lot of work was needed to convert the numbers z and w in Example 3 into polar form, compute their product, and convert back to rectangular form – certainly more work than is required to multiply out $zw = (2\sqrt{3} + 2i)(-1 + i\sqrt{3})$ the old-fashioned way.

However, Theorem 3 pays huge dividends when computing powers of complex numbers. Consider how we computed w^5 above and compare that to using the Binomial Theorem, Theorem ??, to accomplish the same feat by expanding $(-1 + i\sqrt{3})^5$.

Moreover, division is tricky in the best of times, and we saved ourselves a lot of time and effort using Theorem 3 to find and simplify $\frac{z}{w}$ using their polar forms as opposed to starting with $\frac{2\sqrt{3} + 2i}{-1 + i\sqrt{3}}$, rationalizing the denominator, and so forth.


There is geometric reason for studying these polar forms and we would be derelict in our duties if we did not mention the Geometry hidden in Theorem 3.

Take the product rule, for instance. If $z = |z|\operatorname{cis}(\alpha)$ and $w = |w|\operatorname{cis}(\beta)$, the formula $zw = |z||w|\operatorname{cis}(\alpha + \beta)$ can be viewed geometrically as a two step process.


THE POLAR FORM OF COMPLEX NUMBERS

The multiplication of $|z|$ by $|w|$ can be interpreted as magnifying⁹ the distance $|z|$ from z to 0, by the factor $|w|$. Adding the argument of w to the argument of z can be interpreted geometrically as a rotation of β radians counter-clockwise.¹⁰

Focusing on z and w from Example 3, we can arrive at the product zw by plotting z , doubling its distance from 0 (since $|w| = 2$), and rotating $\frac{2\pi}{3}$ radians counter-clockwise. The sequence of diagrams below attempts to describe this process geometrically.



#4
Multiplying z by $|w| = 2$.




#5
Rotating counter-clockwise by $\text{Arg}(w) = \frac{2\pi}{3}$ radians.


Visualizing zw for $z = 4\text{cis}\left(\frac{\pi}{6}\right)$ and $w = 2\text{cis}\left(\frac{2\pi}{3}\right)$.

We may also visualize division similarly. Here, the formula $\frac{z}{w} = \frac{|z|}{|w|}\text{cis}(\alpha - \beta)$ may be interpreted as shrinking¹¹ the distance from 0 to z by the factor $|w|$, followed up by a *clockwise*¹² rotation of β radians.

In the case of z and w from Example 3, we arrive at $\frac{z}{w}$ by first halving the distance from 0 to z , then rotating clockwise $\frac{2\pi}{3}$ radians as shown below.



#6
Dividing z by $|w| = 2$.



#7
Rotating clockwise by $\text{Arg}(w) = \frac{2\pi}{3}$ radians.

Visualizing $\frac{z}{w}$ for $z = 4\text{cis}\left(\frac{\pi}{6}\right)$ and $w = 2\text{cis}\left(\frac{2\pi}{3}\right)$.

Our last goal of the section is to reverse DeMoivre's Theorem to extract roots of complex numbers.

⁹Assuming $|w| > 1$.

¹⁰Assuming $\beta > 0$.

¹¹Again, assuming $|w| > 1$.

¹²Again, assuming $\beta > 0$.

THE POLAR FORM OF COMPLEX NUMBERS

Definition 3. Let z and w be complex numbers. If there is a natural number n such that $w^n = z$, then w is an n^{th} root of z .

Unlike Definition ?? in Section ??, we do not specify one particular *principal* n^{th} root, hence the use of the indefinite article ‘an’ as in ‘an n^{th} root of z ’. Using this definition, both 4 and -4 are square roots of 16, while $\sqrt{16}$ means the principal square root of 16 as in $\sqrt{16} = 4$.

Suppose we wish to find all complex third (cube) roots of 8. Algebraically, we are trying to solve $w^3 = 8$. We know that there is only one *real* solution to this equation, namely $w = \sqrt[3]{8} = 2$, but if we take the time to rewrite this equation as $w^3 - 8 = 0$ and factor, we get $(w - 2)(w^2 + 2w + 4) = 0$.

Solving $w^2 + 2w + 4 = 0$ gives two more cube roots $w = -1 \pm i\sqrt{3}$, for a total of three cube roots of 8. Per Theorem ??, since the degree of $p(w) = w^3 - 8$ is three, there are three complex zeros, counting multiplicity. Since we have found three distinct zeros, we know we have found *all* of the zeros, so there are *exactly three distinct* cube roots of 8.

Let us now solve this same problem using the machinery developed in this section. To do so, we express $z = 8$ in polar form. Since $z = 8$ lies 8 units away on the positive real axis, we get $z = 8\text{cis}(0)$. If we let $w = |w|\text{cis}(\alpha)$ be a polar form of w , the equation $w^3 = 8$ becomes

$$\begin{aligned} w^3 &= 8 \\ (|w|\text{cis}(\alpha))^3 &= 8\text{cis}(0) \\ |w|^3\text{cis}(3\alpha) &= 8\text{cis}(0) \quad \text{DeMoivre's Theorem} \end{aligned}$$

The complex number on the left hand side of the equation corresponds to the point with polar coordinates $(|w|^3, 3\alpha)$, while the complex number on the right hand side corresponds to the point with polar coordinates $(8, 0)$. Since $|w| \geq 0$, so is $|w|^3$, which means $(|w|^3, 3\alpha)$ and $(8, 0)$ are two polar representations corresponding to the same complex number, both with positive r values.

From Section ??, we know $|w|^3 = 8$ and $3\alpha = 0 + 2\pi k$ for integers k . Since $|w|$ is a real number, we solve $|w|^3 = 8$ by extracting the principal cube root to get $|w| = \sqrt[3]{8} = 2$.

As for α , we get $\alpha = \frac{2\pi k}{3}$ for integers k . This produces three distinct points with polar coordinates corresponding to $k = 0, 1$ and 2 : specifically $(2, 0)$, $(2, \frac{2\pi}{3})$ and $(2, \frac{4\pi}{3})$.

The point $(2, 0)$ corresponds to the complex number $w_0 = 2\text{cis}(0)$, the point $(2, \frac{2\pi}{3})$ corresponds to the complex number $w_1 = 2\text{cis}(\frac{2\pi}{3})$, and the point $(2, \frac{4\pi}{3})$ corresponds to the complex number $w_2 = 2\text{cis}(\frac{4\pi}{3})$.

Converting to rectangular form, we find $w_0 = 2$, $w_1 = -1 + i\sqrt{3}$ and $w_2 = -1 - i\sqrt{3}$.

While this process seems a tad more involved than our previous factoring approach, this procedure can be generalized to find, for example, all of the fifth roots of 32. (Try using Chapter ?? techniques on that!)

If we start with a generic complex number in polar form $z = |z|\text{cis}(\theta)$ and solve $w^n = z$ in the same manner as above, we arrive at the following theorem.

Theorem 4. The n^{th} roots of a Complex Number:

Let $z \neq 0$ be a complex number with polar form $z = r\text{cis}(\theta)$. For each natural number n , z has n distinct n^{th} roots, which we denote by $w_0, w_1, \dots, w_{n-1}$, and they are given by the formula

$$w_k = \sqrt[n]{r}\text{cis}\left(\frac{\theta}{n} + \frac{2\pi}{n}k\right)$$

The proof of Theorem 4 breaks into two parts: first, showing that each w_k is an n^{th} root, and second, showing that the set $\{w_k \mid k = 0, 1, \dots, (n-1)\}$ consists of n different complex numbers.

THE POLAR FORM OF COMPLEX NUMBERS

To show w_k is an n^{th} root of z , we use DeMoivre's Theorem to show $(w_k)^n = z$.

$$\begin{aligned}(w_k)^n &= \left(\sqrt[n]{r} \text{cis} \left(\frac{\theta}{n} + \frac{2\pi}{n} k \right) \right)^n \\&= (\sqrt[n]{r})^n \text{cis} \left(n \cdot \left[\frac{\theta}{n} + \frac{2\pi}{n} k \right] \right) \quad \text{DeMoivre's Theorem} \\&= r \text{cis} (\theta + 2\pi k)\end{aligned}$$

Since k is a whole number, $\cos(\theta + 2\pi k) = \cos(\theta)$ and $\sin(\theta + 2\pi k) = \sin(\theta)$. Hence, it follows that $\text{cis}(\theta + 2\pi k) = \text{cis}(\theta)$, so $(w_k)^n = r \text{cis}(\theta) = z$, as required.

To show that the formula in Theorem 4 generates n distinct numbers, we assume $n \geq 2$ (or else there is nothing to prove) and note that the modulus of each of the w_k is the same, namely $\sqrt[n]{r}$.

Therefore, the only way any two of these polar forms correspond to the same number is if their arguments are coterminal – that is, if the arguments differ by an integer multiple of 2π .

Suppose k and j are whole numbers between 0 and $(n-1)$, inclusive, with $k \neq j$. Since k and j are different, let's assume for the sake of argument that $k > j$. Then $\left(\frac{\theta}{n} + \frac{2\pi}{n} k \right) - \left(\frac{\theta}{n} + \frac{2\pi}{n} j \right) = 2\pi \left(\frac{k-j}{n} \right)$.

For $2\pi \left(\frac{k-j}{n} \right)$ to be an integer multiple of 2π , $(k-j)$ must be a multiple of n . But because of the restrictions on k and j , $0 < k-j \leq n-1$. (Think this through.) Hence, $(k-j)$ is a positive number less than n , so it cannot be a multiple of n .

As a result, w_k and w_j are different complex numbers, and we are done. By Theorem ??, we know there at most n distinct solutions to $w^n = z$, and we have just found all n of them.

We illustrate Theorem 4 in the next example.

Example 4. Use Theorem 4 to find the following:

1. both square roots of $z = -2 + 2i\sqrt{3}$
2. the four fourth roots of $z = -16$
3. the three cube roots of $z = \sqrt{2} + i\sqrt{2}$
4. the five fifth roots of $z = 1$.

Solution.

1. We start by writing $z = -2 + 2i\sqrt{3}$ in polar form as $z = 4 \text{cis} \left(\frac{2\pi}{3} \right)$. Since we are looking for *square* roots, $n = 2$. In keeping with the notation used in Theorem 4 we will call these roots w_0 and w_1 , in keeping with the notation suggested there.

Identifying $r = 4$, $\theta = \frac{2\pi}{3}$, Theorem 4 gives one root as $w_0 = \sqrt{4} \text{cis} \left(\frac{(2\pi/3)}{2} + \frac{2\pi}{2}(0) \right) = 2 \text{cis} \left(\frac{\pi}{3} \right)$ and the other root as $w_1 = \sqrt{4} \text{cis} \left(\frac{(2\pi/3)}{2} + \frac{2\pi}{2}(1) \right) = 2 \text{cis} \left(\frac{4\pi}{3} \right)$.

Though not asked to do so, we can easily convert each of w_0 and w_1 to rectangular form: $w_0 = 1 + i\sqrt{3}$ and $w_1 = -1 - i\sqrt{3}$. We can check our answers by showing $w_0^2 = -2 + 2i\sqrt{3}$ and $w_1^2 = -2 + 2i\sqrt{3}$.

2. Proceeding as above, we begin by converting z to polar form: $z = -16 = 16 \text{cis}(\pi)$. Here, $n = 4$, so Theorem 4 guarantees us *four* fourth roots.

THE POLAR FORM OF COMPLEX NUMBERS

Identifying $r = 16$, $\theta = \pi$ and $n = 4$, Theorem 4 gives us: $w_0 = \sqrt[4]{16}\text{cis}\left(\frac{\pi}{4} + \frac{2\pi}{4}(0)\right) = 2\text{cis}\left(\frac{\pi}{4}\right)$, $w_1 = \sqrt[4]{16}\text{cis}\left(\frac{\pi}{4} + \frac{2\pi}{4}(1)\right) = 2\text{cis}\left(\frac{3\pi}{4}\right)$, $w_2 = \sqrt[4]{16}\text{cis}\left(\frac{\pi}{4} + \frac{2\pi}{4}(2)\right) = 2\text{cis}\left(\frac{5\pi}{4}\right)$ and last, but not least, $w_3 = \sqrt[4]{16}\text{cis}\left(\frac{\pi}{4} + \frac{2\pi}{4}(3)\right) = 2\text{cis}\left(\frac{7\pi}{4}\right)$.

Once again, we can conveniently convert our answers to rectangular form. We get: $w_0 = \sqrt{2} + i\sqrt{2}$, $w_1 = -\sqrt{2} + i\sqrt{2}$, $w_2 = -\sqrt{2} - i\sqrt{2}$ and $w_3 = \sqrt{2} - i\sqrt{2}$. We invite the reader to check our answers algebraically by showing $w_0^4 = w_1^4 = w_2^4 = w_3^4 = -16$.

3. For $z = \sqrt{2} + i\sqrt{2}$, we have $z = 2\text{cis}\left(\frac{\pi}{4}\right)$. With $r = 2$, $\theta = \frac{\pi}{4}$ and $n = 3$ the usual computations yield $w_0 = \sqrt[3]{2}\text{cis}\left(\frac{\pi}{12}\right)$, $w_1 = \sqrt[3]{2}\text{cis}\left(\frac{9\pi}{12}\right) = \sqrt[3]{2}\text{cis}\left(\frac{3\pi}{4}\right)$ and $w_2 = \sqrt[3]{2}\text{cis}\left(\frac{17\pi}{12}\right)$.

To convert our answers to rectangular form requires the use of either the Sum and Difference Identities in Theorem ?? or the Half-Angle Identities in Theorem ?? to evaluate w_0 and w_2 . Since we are not explicitly told to do so, we leave this as a good, but messy, exercise.

4. To find the five fifth roots of 1, we write $1 = 1\text{cis}(0)$. We have $r = 1$, $\theta = 0$ and $n = 5$. Since $\sqrt[5]{1} = 1$, the roots are $w_0 = \text{cis}(0) = 1$, $w_1 = \text{cis}\left(\frac{2\pi}{5}\right)$, $w_2 = \text{cis}\left(\frac{4\pi}{5}\right)$, $w_3 = \text{cis}\left(\frac{6\pi}{5}\right)$ and $w_4 = \text{cis}\left(\frac{8\pi}{5}\right)$.


The situation here is even graver than in the previous example, since we have not developed any identities to help us determine the cosine or sine of $\frac{2\pi}{5}$. At this stage, we could approximate our answers using a calculator, and we leave this as an exercise. ■

Having done some computations with Theorem 4, it's time to take a step back to look at things geometrically.

Essentially, Theorem 4 says that to find the n^{th} roots of a complex number, we first take the n^{th} root of the modulus and divide the argument by n . This gives the first root w_0 .

Each successive root is found by adding $\frac{2\pi}{n}$ to the argument, which amounts to rotating w_0 by $\frac{2\pi}{n}$ radians. The result of these actions produces n roots, spaced equally around the complex plane.

As an example of this, we plot our answers to number 2 in Example 4 below.



The four fourth roots of $z = -16$ equally spaced $\frac{2\pi}{4} = \frac{\pi}{2}$ around the plane.

We have only glimpsed at the beauty of the complex numbers in this section. The complex plane is without a doubt one of the most important mathematical constructs ever devised. Coupled with Calculus, it is the venue for incredibly important Science and Engineering applications.¹³

¹³For more on this, see the beautifully written epilogue to Section ?? found on page ??.